

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &  
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA  
AMFU, OFRS, NAGANAHALLI,  
MYSURU – 570003**



**Date: 01-08-2025**

**AGRO-ADVISORY BULLETIN FOR CHAMARAJANAGARA DISTRICT**

**Issued jointly by, UAS, Bengaluru & Indian Meteorological Department**

**Past Weather Data**

<b>Parameter</b>	<b>29.07.2025</b>	<b>30.07.2025</b>	<b>31.07.2025</b>	<b>01.08.2025</b>
<b>Rainfall (mm)</b>	-	-	-	-
<b>Max. Temp. (°C)</b>	-	-	-	-
<b>Min. Temp. (°C)</b>	-	-	-	-
<b>Sky condition (Octas)</b>	-	-	-	-
<b>Relative humidity (%) 0830 hours</b>	-	-	-	-
<b>Relative humidity (%) 1730 hours</b>	-	-	-	-
<b>Wind Speed (km/h)</b>	-	-	-	-
<b>Wind Direction</b>	-	-	-	-

**Weather forecast for the next five days (From 02-08-2025 to 06-08-2025)**

<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	1	2	4	6	15
<b>Max. Temp. (°C)</b>	29	30	29	29	29
<b>Min.Temp. (°C)</b>	20	20	20	20	20
<b>Sky condition (Octas)</b>	6	5	6	6	7
<b>Relative humidity (%) 0830 hours</b>	85	85	87	89	91
<b>Relative humidity (%) 1730 hours</b>	65	64	62	63	62
<b>Wind Speed (kmph)</b>	12	12	12	12	12
<b>Wind Direction</b>	272	279	279	277	276

\*GW - Gusty wind, TSH – Thunderstorm

**Forecast Summary**

As forecast received from IMD, partially cloudy sky with **light rainfall** may be expected from 02.08.2025 to 06.08.2025 in Chamarajanagara district. The day temperature is expected to be 29-30°C & night temperature is expected 20°C. The relative humidity in the morning hours is expected to be 85-91% & afternoon relative humidity is expected to be in the range of 62-65%. Wind speed expected to be 12 km/ hr.

**General Advisory:**

- ✓ **Wind Warning:** All crops are vulnerable to lodging or physical damage from gusty winds—take preventive structural/mechanical measures.
- ✓ **Disease Precaution:** High relative humidity (especially mornings) favors fungal/bacterial diseases—monitor crops closely.
- ✓ **Field Operations:** Schedule intercultural and pesticide applications during rain-free, low-wind periods.

**SMS Advisory:**

Rain and strong winds expected; support crops, ensure drainage, delay paddy transplanting during peak winds, and protect tomato and chilli from disease.

**Crops and Suitable Varieties for Sowing in July****Paddy (Rice)**

- **General Sowing:** MSN-99
- **Late June (3rd-4th week):** BR-2655, Jaya
- **Aerobic Paddy:** Daksha (KMP-175), Onnasiri (MAS-26), Sharada (MAS-946), Anagha (ARB-6), Rashi

**Ragi (Finger Millet):** Indaf-8, MR-1, MR-6, L-5, KMR-301, GPU-45, KMR-316

**Maize (Shaktiman Maize):** CSH-5, CSH-9

**Pearl Millet (Musuku Jola):** Hema, Nithyashree, MAH-14-5, MAH-14-138, MAH-15-84

**Popcorn:** Amber

**Bajra (Pearl Millet):** WCC-75, PHVB-910

- **Late June (3rd-4th week):** ICTP-8203, SMV-221, MH-946 (GHB-558)

**Foxtail Millet (Navane):** PS-4, SIA-326, GPUF-3

**Horsegram (Haraka):** RBK-155, GPUK-3

**Little Millet (Same):** CO-2, OLM-203, GPUL-6, GPUL-11

**Proso Millet (Oodalu):** DHBM-93-3

**Kodo Millet (Korale):** GPUBT-2

**Red Gram (Pigeon Pea / Togari):** BRG-1, BRG-2, BRG-3, BRG-4, BRG-5, ICP-7035, TTB-7, Hyderabad-3C

**Groundnut (Nelekadale):** GKVK-5, KCG-6 (Chintamani-6), GPBD-4, ICGV-91114, JL-24, TMV-2, GKVK-27

**Sunflower:** KBSH-41, 90, 44, 53, 78, 85

**Soybean (Soya Avare):** MAUS-2 (Pooja), Karune (vegetable soybean), KBS-23

**Cotton:** DCS-9 (Jyoti), 48-1 (Jwala)

- **Hybrid Varieties:** DCH-177, GCH-4, 30.N.23-66

**Sesame (Hucchellu) - 4th week of June:** KBN-1, No.71, KBN-2, 52: 2.2-62175, 2.2-86032, 2.2.A.2-18061

**Horticultural Crops**

- **Vegetables:** Tomato, Potato, Brinjal, Chilli, Cabbage, Radish, Knol-khol, Carrot, Onion
- **Fruits:** Mango, Banana, Lime, Guava, Sweet lime, Sapota, Papaya, Pomegranate, Pineapple, Cashew
- **Spices:** Garlic, Ginger, Turmeric, Cardamom

**Fodder Crops**

- **Sorghum (Fodder Maize):** JS-SF-3
- **Cowpea (Fodder):** MF-09-01, MF-08-14, 2-09-03
- **Napier-Bajra Hybrid Grass:** PBN-342

Recommendations to the farmers:-		
Weather based advisory		
Crop	Stage	Weather-Based Advisory
Paddy	Transplanting	<ul style="list-style-type: none"> <li>- Avoid transplanting during gusty wind hours; secure young seedlings.</li> <li>- Maintain optimum water level (2–3 cm) in field.</li> <li>- Apply recommended basal fertilizer post-transplanting.</li> </ul>
Maize	Vegetative	<ul style="list-style-type: none"> <li>- Provide staking/support against gusty winds.</li> <li>- Monitor for stem borer infestation due to high humidity.</li> <li>- Ensure proper drainage.</li> </ul>
Tomato	Fruit formation	<ul style="list-style-type: none"> <li>- Prevent fungal diseases (blight) using recommended fungicides due to high humidity &amp; rainfall.</li> <li>- Avoid water stagnation.</li> </ul>
Chilli	Fruit formation	<ul style="list-style-type: none"> <li>- Spray preventive fungicides for anthracnose and fruit rot.</li> <li>- Harvest mature fruits before rainfall if possible.</li> </ul>
Banana	Fruit development	<ul style="list-style-type: none"> <li>- Provide propping to prevent lodging from gusty winds.</li> <li>- Maintain drainage.</li> <li>- Apply potassium-rich fertilizer for fruit quality.</li> </ul>
Black gram, Green gram, Red gram, Field bean, Cowpea	Vegetative	<ul style="list-style-type: none"> <li>- Check for pest/disease incidence, especially leaf spots and aphids.</li> <li>- Avoid foliar sprays during rainfall.</li> <li>- Ensure soil is not waterlogged.</li> </ul>
Sugarcane	Vegetative	<ul style="list-style-type: none"> <li>- Ensure sufficient drainage.</li> <li>- Consider earthing-up operations.</li> <li>- Apply recommended fertilizer dose post-rain.</li> </ul>
Vegetable crops (general)	Various stages	<ul style="list-style-type: none"> <li>- Protect from fungal infections with prophylactic sprays.</li> <li>- Provide support where needed.</li> <li>- Harvest mature vegetables before expected rainfall.</li> </ul>

Sector	Advisory
Livestock	<ul style="list-style-type: none"> <li>- Provide clean, dry shelter to protect from rain and wind.</li> <li>- Maintain hygiene to prevent hoof and skin infections.</li> <li>- Ensure adequate clean water and balanced nutrition.</li> </ul>
Poultry	<ul style="list-style-type: none"> <li>- Keep sheds well-ventilated but protected from rain and gusty winds.</li> <li>- Maintain dry litter to avoid disease.</li> <li>- Provide vitamin supplements to boost immunity.</li> </ul>
Sericulture	<ul style="list-style-type: none"> <li>- Maintain optimum temperature and humidity in rearing houses.</li> <li>- Ensure mulberry leaves are not wet during feeding.</li> <li>- Improve ventilation to prevent fungal diseases in silkworms.</li> </ul>

Recommendation to farmers		
Crop specific advisory:		
Crop	Stage	Advisory
Maize fall army worm	Tasseling stage	<ul style="list-style-type: none"> <li>✓ Handpick and destroy egg masses and larvae.</li> <li>✓ Use predators like <i>Trichogramma pretiosum</i> or parasitoids like <i>Telenomus remus</i>.</li> <li>✓ Apply <i>Metarhizium anisopliae</i> or <i>Beauveria bassiana</i>.</li> <li>✓ Spray Chlorantraniliprole 18.5% SC @ 0.4 ml/l or Emamectin benzoate 5% SG @ 0.4 g/l. Avoid excessive nitrogen application.</li> </ul>

<b>Coconut rugose whitefly</b>	Vegetative stage	<ul style="list-style-type: none"> <li>✓ Prune and burn infested leaves.</li> <li>✓ Release <i>Encarsia guadeloupae</i> parasitoids. Conserve natural predators like ladybird beetles (<i>Cryptolaemus montrouzieri</i>).</li> <li>✓ Spray Neem oil 1% or use Acephate 75 SP @ 1 g/l as a spot application if infestation is severe.</li> </ul>
<b>Chilli leaf curl virus</b>	Fruit formation stage	<ul style="list-style-type: none"> <li>✓ Use virus-free seeds and resistant varieties. Maintain proper spacing and avoid overlapping.</li> <li>✓ Remove and destroy infected plants. Use yellow sticky traps to monitor whitefly populations.</li> <li>✓ Spray Imidacloprid 17.8% SL @ 0.5 ml/l or Thiamethoxam 25 WG @ 0.3 g/l.</li> </ul>
<b>Tomato Early and late blight of tomato</b>	Flowering	<p>For late blight of tomato 15 days prior to transplanting Trichoderma and Pseudomonas enriched compost may be incorporated to the soil. For early blight control spray 2.0 g. Mancozeb 75 WP</p> <p>OR</p> <p>2.0 g. Maneb</p> <p>OR</p> <p>2.0 g. Metalaxyl- MZ 72WP.</p> <p>OR</p> <p>2.0 g. Dimethomorph + polyram/lit. water.</p> <p>For control of late blight spray 2.0 g. Metalaxyl - MZ 72WP.</p> <p>OR</p> <p>2.0 g. Fosetyl al 80 WP</p> <p>OR</p> <p>2.0 g. Dimethomorph + polyram in a lit. water, 5 weeks after transplanting. Repeat the spray 7th, 9th and 11th weeks after transplanting. 200- 250 lit. spray solution required/acre/spray.</p>

**Block level weather forecast (From 02-08-2025 to 06-08-2025)**

**Chamarajanagara**

<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	2.1	4.7	7.8	8.3	12.4
<b>Max. temp (°C)</b>	29	28.7	28.5	27.2	25.4
<b>Min.Temp (°C)</b>	20.2	20.2	20.2	20.1	20.1
<b>Sky condition (Octas)</b>	8	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	92.8	91.6	92.7	93.6	91.1
<b>Relative humidity (%) 1730 hours</b>	60.1	60.6	52.4	57.7	67.6
<b>Wind Speed (kmph)</b>	8.9	8	7.6	8	7.8
<b>Wind Direction</b>	284	288.4	289.3	280.3	283.4

**Gundlupete**

<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	1.8	4.8	6.1	7.5	13.2
<b>Max. temp (°C)</b>	27.9	27.5	27.6	26.4	25
<b>Min.Temp (°C)</b>	19.9	20.2	20.1	19.9	19.8
<b>Sky condition (Octas)</b>	7	8	8	8	8

<b>Relative humidity (%) 0830 hours</b>	91.5	91.8	91.2	92.5	92.3
<b>Relative humidity (%) 1730 hours</b>	65.4	61.8	56.3	60.8	72.2
<b>Wind Speed (kmph)</b>	9.7	9.4	8.3	9.8	8.3
<b>Wind Direction</b>	360	274.4	275	263.7	360

### Kollegala

<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	3.8	7.1	12.6	12.8	10.1
<b>Max. temp (°C)</b>	29.9	29.4	29.5	28	26.5
<b>Min.Temp (°C)</b>	21	21	21.1	21	20.7
<b>Sky condition (Octas)</b>	7	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	93.4	93.6	93.5	92.9	92.1
<b>Relative humidity (%) 1730 hours</b>	60.6	63.2	51.6	58.4	68.6
<b>Wind Speed (kmph)</b>	8.4	7.1	7.2	8	7.2
<b>Wind Direction</b>	244.5	255.2	252.5	243.4	252.5

### Yelandur

<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	4.1	9	12.9	13.7	11.2
<b>Max. temp (°C)</b>	29.6	29	29.1	27.7	26.1
<b>Min.Temp (°C)</b>	20.8	20.8	20.8	20.7	20.5
<b>Sky condition (Octas)</b>	8	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	93.5	93.1	93.8	94	92.4
<b>Relative humidity (%) 1730 hours</b>	61.2	62.4	52.7	59.8	69.1
<b>Wind Speed (kmph)</b>	7.6	6.6	6.6	7.3	6.6
<b>Wind Direction</b>	250.7	260.5	257.5	249.8	260.5

### Hanur


<b>Parameter</b>	<b>02.08.2025</b>	<b>03.08.2025</b>	<b>04.08.2025</b>	<b>05.08.2025</b>	<b>06.08.2025</b>
<b>Rainfall (mm)</b>	3.4	8	15.4	13.4	11.4
<b>Max. temp (°C)</b>	28.7	28.4	27.8	26	24.9
<b>Min.Temp (°C)</b>	20	20.2	20.1	20	19.9
<b>Sky condition (Octas)</b>	8	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	94.2	92.9	96	93	93
<b>Relative humidity (%) 1730 hours</b>	61.1	61.6	54.5	62.4	69.3
<b>Wind Speed (kmph)</b>	9	8.3	7.6	7.9	7.6
<b>Wind Direction</b>	267.7	272.5	360	267.4	360

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)


For any information farmers can contact **Dr.Umashankar Kumar, N.**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Technical officer over phone No. 0821-259126/ 9535345814.

**AMFU of IMD,  
Naganahalli, Mysuru**

वास्तविकवर्षा तथा विस्तारित अवधि पूर्वानुमान  
**Realized Rainfall and Extended Range Forecast**  
 (वर्षा और तापमान)  
 (Rainfall and Temperature)








- Week1(01.08.2025 to 07.08.2025):** Rainfall is likely to be above normal over many parts of North West India, East India and Tamil Nadu. However, it is likely to be below normal over West India, coastal Karnataka, Telangana, many parts of Central India, Nagaland Manipur Mizoram & Tripura, Coastal Andhra Pradesh and Odisha.
- Week 2 (08.08.2025 to 14.08.2025):** Rainfall is likely to be above normal over many parts of South India, East India, some parts of North East India, East Uttar Pradesh, Madhya Maharashtra, Haryana and adjoining parts of Rajasthan. However, it is likely to be below normal over Central India, Gujarat, Konkan-Goa, Jammu & Kashmir, Punjab, Himachal Pradesh and Uttarakhand.

### Maximum and Minimum temperature anomaly ( $^{\circ}$ C) forecast for the next 2 weeks (IC- 30<sup>th</sup> July, 2025) (01<sup>st</sup> to 14<sup>th</sup> August, 2025)




### Maximum Temperature (Tmax)

- Week 1 (01.08.2025 to 07.08.2025):** Maximum temperature is likely to be above normal over Odisha, Chhattisgarh, Karnataka, Madhya Maharashtra, some parts of Madhya Pradesh and Coastal Andhra Pradesh.
- Week 2 (08.08.2025 to 14.08.2025):** Maximum temperature is likely to be above normal over Central India, West India, Karnataka, Himachal Pradesh, parts of Telangana and Coastal Andhra Pradesh.

### MME forecast Tmin anomaly (Deg C)

(Week1: 01Aug–07Aug)

(Week2: 08Aug–14Aug)



#### Minimum Temperature (Tmin)

- **Week 1 (01.08.2025 to 07.08.2025):** Minimum temperature is likely to be above normal over Karnataka, Chhattisgarh and Himachal Pradesh.
- **Week 2 (08.08.2025 to 14.08.2025):** Minimum temperature is likely to be above normal over many parts of Central India, North West India, Maharashtra and Karnataka.