

**OF AGRICULTURAL SCIENCES, BENGALURU &
INDIA METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA AMFU,
AICRP- Agrometeorology, UAS,GKVK
Bengaluru – 560 065**



Date: 13-02-2026

**AGRO-ADVISORY BULLETIN FOR Bengaluru South DISTRICT
Issued jointly by UAS, Bangalore & Indian Meteorological Department**

Past Weather Data (09-02-2026 to 13-02-2026)

	09.02.2026	10.02.2026	11.02.2026	12.02.2026	13.02.2026
Rainfall (mm)	-	-	-	-	-
Max. Temp. (°C)	-	-	-	-	-
Min. Temp. (°C)	-	-	-	-	-
Sky condition (Octas)	-	-	-	-	-
Relative humidity (%) 0830 hours	-	-	-	-	-
Relative humidity (%) 1730 hours	-	-	-	-	-
Wind Speed (km/h)	-	-	-	-	-
Wind Direction	-	-	-	-	-



Weather forecast for the next five days (From 14-02-2026 to 18-02-2026)

Parameter	14.02.2026	15.02.2026	16.02.2026	17.02.2026	18.02.2026
Rainfall (mm)	0	0	0	0	0
Max. Temp. (°C)	31	31	30	30	31
Min. Temp. (°C)	17	17	17	16	17
Sky condition (Octas)	3	2	3	4	4
Relative humidity (%) 0830 hours	66	65	67	64	38
Relative humidity (%) 1730 hours	33	35	34	35	34
Wind Speed (kmph)	4	6	6	6	4
Wind Direction	76	88	88	98	97

Forecast Summary

As forecast received from IMD, cloudy sky with **No rain** expected 14-02-2026 to 18-02-2026 in Bengaluru South District. The day temperature is expected to be 30.0-31.0°C and night temperature is expected to be 16.0-17.0°C. The relative humidity in the morning hours is expected to be 38-67 % and afternoon relative humidity is expected to be in the range of 33-35 %, Wind speed is expected to be 4-6 km/hr.

SMS Advisory

Avoid pruning, fertilizer application and growth regulator sprays during cold wave conditions. Cloudy weather, cool nights may increase fungal disease risk—monitor crops closely and ensure fields have good drainage.



Recommendations to the farmers: -

Crop	Pest/Disease	Damage symptoms	Control measures
------	--------------	-----------------	------------------

General Advisory:

Field Crops


1. Right time for harvesting, drying, cleaning and storage of Rabi crops.
2. Apply **neem leaves/neem powder** in grain bags as a natural repellent.
3. For long-term storage of pulses, store with **tri-sodium phosphate (TSP) treated** gunny sacks to reduce bruchid attack.

Vegetables & Horticulture

1. Watch for fruit borer and shoot borer in fruit development stage.
2. Spray the chemicals early morning and late evening for better pest and disease control.

Livestock & Poultry

1. Give dry fodder and provide shelter to animals in evening higher humidity.
2. Maintain hygiene in sheds to prevent infections.



Crop	Stage	Pest and Diseases	Weather-Based Agromet Advisory
Ragi	Post harvest		Dry the harvested ear heads on clean tarpaulins until grain moisture reaches about 12%.
Redgram	Post harvest		Dry the harvested pods in 3-4 days and separate the seeds and store in cool places The grain moisture reaches about 14%.
Cowpea	Post harvest stage		Harvest the matured pod and dry in clean tarpaulins until grain moisture reaches about 15%.
Field bean	Post harvest stage		Harvest the matured pod and dry in clean tarpaulins until grain moisture reaches about 15%.
Chilli	Fruit development stage	Thrips	Spray Carbendazim 1 g/l of water or Copper oxychloride 2.5 g/l of water for anthracnose. For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water. Avoid water stagnation in crop field.
Mango	Marble stage	Powdery mildew and Green leaf hopper	To control of green leaf hopper in Mango spray Imidacloprid 17.8 SL @ 0.3 ml /l of water To control of powdery mildew in Mango spray wettable sulphur @ 3 g /l of water in leaf and flower parts of affected parts.
Arecanut		Spindle bug	Moist conditions may cause infestation of spindle bug and inflorescence dieback . To control inflorescence dieback spray Copper oxychloride 3 g/litre on bunches and crown region. To control spindle bug spray Dimethoate 30 EC @ 2



			ml/litre of water. Spray on spindle leaves and crown region using a hand sprayer
--	--	--	--

Livestock, Poultry, and Sericulture Advisory	
Sector	Weather-Based Advisory
Livestock	<ol style="list-style-type: none"> 1. Provide dry and clean shelter; avoid animals standing in wet areas. 2. Provide ample clean drinking water. 3. Monitor for tick and mite infestations; use approved acaricides if needed. 4. Provide balanced feed and mineral supplements. 5. Minimum temperatures cause cold stress in young calves/kids. Provide bedding (dry straw) and night shelter to reduce cold exposure.
Sericulture	<p>Humid and rainy conditions increase grasserie, flacherie and fungal diseases</p> <ol style="list-style-type: none"> 1. Maintain proper rearing house hygiene, clean and disinfect trays. 2. The recommendation of farmers closes the windows with tarpaulins sheet during night hours to maintain optimum room temperature. 3. Avoid wet or damp mulberry leaves, use air-dried or well-drained leaves.
Poultry	<ol style="list-style-type: none"> 1. Cool early mornings need brooder temperature maintenance for chicks. 2. Maintain poultry shed dryness; use lime powder to reduce moisture. 3. Provide electrolytes + vitamins in water for immunity. 4. Cool, humid mornings favor: CRD (Chronic Respiratory Disease), Coccidiosis, Colibacillosis- Follow routine vaccinations strictly (Ranikhet, IBD). 5. Maintain optimum lighting schedule to support winter egg production.

Block level weather forecast (From 14-02-2026 to 18-02-2026)					
CHANNAPATTANA BLOCK					
Parameter	14.02.2026	15.02.2026	16.02.2026	17.02.2026	18.02.2026
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	33	33	33	33	33
Min.Temp (°C)	16	16	16	17	17
Sky condition (Octas)	2	3	2	3	4
Relative humidity (%) 0830 hours	83	74	68	66	71
Relative humidity (%) 1730 hours	21	16	13	13	18
Wind Speed (kmph)	7	8	6	6	7
Wind Direction	79	68	76	83	90

KANAKAPURA BLOCK					
Parameter	14.02.2026	15.02.2026	16.02.2026	17.02.2026	18.02.2026
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	32	33	33	33	32

Min.Temp (°C)	16	16	16	17	17
Sky condition (Octas)	3	3	2	3	3
Relative humidity (%) 0830 hours	85	79	73	67	71
Relative humidity (%) 1730 hours	23	17	13	15	19
Wind Speed (kmph)	5	6	5	6	6
Wind Direction	90	83	82	94	97

MAGADI BLOCK

Parameter	14.02.2026	15.02.2026	16.02.2026	17.02.2026	18.02.2026
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	32	32	32	32	32
Min.Temp (°C)	16	16	16	16	16
Sky condition (Octas)	2	2	3	3	4
Relative humidity (%) 0830 hours	73	70	64	64	68
Relative humidity (%) 1730 hours	18	16	14	13	18
Wind Speed (kmph)	8	9	8	10	10
Wind Direction	85	81	95	101	112

RAMANAGARA BLOCK


Parameter	14.02.2026	15.02.2026	16.02.2026	17.02.2026	18.02.2026
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	33	33	33	33	33
Min.Temp (°C)	16	16	16	17	17
Sky condition (Octas)	2	3	2	3	4
Relative humidity (%) 0830 hours	81	73	67	65	71
Relative humidity (%) 1730 hours	20	16	13	13	18
Wind Speed (kmph)	8	8	7	7	7
Wind Direction	82	70	78	87	90

- Download “DAMINI” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “MAUSAM” APP for location specific forecast & warning & “MEGHDOOT” APP for Agromet advisory
- This information is available in the website: mausam.imd.gov.in

For any information farmers can contact **Dr. M. N. Thimmegowda**, Professor & Head/
Mr. L. Nagesha, Technical officer over phone No. **9741109702/ 9008454142**


**AMFU of IMD,
AICRP-AM, Bengaluru**

वास्तविक वर्षा तथा विस्तारित अवधि पूर्वानुमान
Realized Rainfall and Extended Range Forecast
 (वर्षा और तापमान)
(Rainfall and Temperature)



Extended Range Forecast System

Rainfall forecast maps for the next 2 weeks (IC- 21st January,2026) (23rd January to 05th February, 2026)




- **Week 1 (23.01.2025 to 29.01.2026):** Rainfall associated with Western Disturbance is likely to be above normal over many parts of Northwest India.
- **Week 2 (30.01.2025 to 05.02.2026):** Rainfall activity is likely over Jammu & Kashmir, Himachal Pradesh and Arunachal Pradesh.

Maximum and Minimum temperature anomaly (°C) forecast for the next 2 weeks (IC- 21st January,2026) (23rd January to 05th February,2026)

MME forecast Tmax anomaly (Deg C)

(Week1: 23Jan–29Jan)

(Week2: 30Jan–05Feb)




Maximum Temperature (Tmax)

- **Week 1 (23.01.2025 to 29.01.2026):** Maximum temperature is likely to be above normal over Jammu & Kashmir, Chhattisgarh, Odisha, Arunachal Pradesh, Assam, North Coastal Andhra Pradesh and some parts of Uttarakhand. However, it is likely to be below normal over Punjab, Haryana, Rajasthan, West India and many parts of Central & South India.
- **Week 2 (30.01.2025 to 05.02.2026):** Maximum temperature is likely to be above normal over North West India, Chhattisgarh, Odisha, Arunachal Pradesh, Assam and parts of Coastal Andhra Pradesh & South Karnataka. However, it is likely to be below normal over North Karnataka, Telangana, Rayalaseema and many parts of Central & West India.

MME forecast Tmin anomaly (Deg C)

(Week1: 23Jan–29Jan)

(Week2: 30Jan–05Feb)



Minimum Temperature (Tmin)

- **Week 1 (23.01.2025 to 29.01.2026):** Minimum temperature is likely to be below normal over Chhattisgarh, Vidarbha and many parts of North West, East & South India. However,

it is likely to be above normal over North East India, many parts of Jammu & Kashmir and some parts of West Madhya Pradesh, Madhya Maharashtra & South Karnataka.

- **Week 2 (30.01.2025 to 05.02.2026):** Minimum temperature is likely to be below normal over East India, West Rajasthan, Chhattisgarh, Telangana, Kerala and Tamil Nadu. However, it is likely to be above normal over many parts of North West India, North East India, Gujarat, Madhya Pradesh, Madhya Maharashtra and South Karnataka.