


**OF AGRICULTURAL SCIENCES, BENGALURU &
INDIA METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA AMFU,
AICRP- Agrometeorology, UAS,GKVK
Bengaluru – 560 065**



Date: 05-05-2026

**AGRO-ADVISORY BULLETIN FOR Bengaluru South DISTRICT
Issued jointly by UAS, Bangalore & Indian Meteorological Department**

Past Weather Data (01-05-2026 to 05-05-2026)

	01.05.2026	02.05.2026	03.05.2026	04.05.2026	05.05.2026
Rainfall (mm)	-	-	-	-	-
Max. Temp. (°C)	-	-	-	-	-
Min. Temp. (°C)	-	-	-	-	-
Sky condition (Octas)	-	-	-	-	-
Relative humidity (%) 0830 hours	-	-	-	-	-
Relative humidity (%) 1730 hours	-	-	-	-	-
Wind Speed (km/h)	-	-	-	-	-
Wind Direction	-	-	-	-	-



Weather forecast for the next five days (From 06-05-2026 to 10-05-2026)

Parameter	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
Rainfall (mm)	2	3	4	5	5
Max. Temp. (°C)	33	33	34	33	33
Min. Temp. (°C)	23	23	24	23	23
Sky condition (Octas)	5	5	5	6	6
Relative humidity (%) 0830 hours	72	74	73	72	71
Relative humidity (%) 1730 hours	45	46	47	45	44
Wind Speed (kmph)	12	10	6	6	6
Wind Direction	234	229	256	262	246

Forecast Summary

As forecast received from IMD, clear sky with **Light rainfall with gusty wind expected during 06-05-2026 to 10-06-2026** in Bengaluru South District. The day temperature is expected to be 33.0-34.0°C and night temperature is expected to be 23.0-24.0°C. The relative humidity in the morning hours is expected to be 71-74 % and afternoon relative humidity is expected to be in the range of 45-47 %, Wind speed is expected to be 6-12 km/hr.

SMS Advisory

Due to increase in temperature day by day promotes fruit drop is observed in mango orchard. Provide one or two irrigations at marble, fruit development and maturity stage of crop



Recommendations to the farmers: -

Crop	Pest/Disease	Damage symptoms	Control measures
------	--------------	-----------------	------------------


General Advisory:

Field Crops

1. **Summer Ploughing:** Perform deep ploughing immediately after the previous harvest to expose the soil to the sun. This kills hibernating pests, pathogens, and weed seeds.
2. **Soil Testing:** Collect soil samples before applying any amendments. This ensures precise nutrient application
3. **In-situ Moisture Conservation:** Repair and strengthen field bunds to prevent runoff and ensure every drop of early monsoon rain is retained in the soil.

Vegetables & Horticulture

1. Watch for fruit borer and shoot borer in fruit development and maturity stage.
2. Spray the chemicals early morning and late evening for better pest and disease control.



Crop	Stage	Pest and Diseases	Weather-Based Agromet Advisory
Pre Kharif land preparation			Apply well-decomposed Farmyard Manure (FYM) or compost at 10–12 tons/ha at least 15–20 days before sowing to improve soil structure and water-holding capacity. Tank Silt: Application of tank silt @ 20-25 tonne/acre of land to improve water holding capacity of soil. Green Manuring: If early showers occur, sow green manure crops like Sunnhemp or Dhaincha and incorporate them into the soil after 45 days to boost nitrogen levels.
French beans	Flowering	Mites	Avoid water stress during flowering and frequent irrigation is required for better growth. To improve pod setting in summer, a foliar spray of Solubor (Boron) @ 1g/liter is recommended. This strengthens the pollen tube and reduces flower drop. Spray Wettable Sulphur 2 g/L
Chilli	Fruit development stage	Thrips	For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water. Apply recommended dose of potash to improve fruit quality and pungency. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear.
Ridge guard	Flowering and fruit formation stage	Fruit Fly	Install Cue-lure pheromone traps (10/acre). Wrap young fruits in paper/cloth bags. Irrigate at 3–4 day intervals in summer.
Tomato	Flowering	Whitefly / Aphids	Flower Drop: When temperatures >32 °C during the day or 24 °C at night, pollen becomes sterile and flowers fall. Spray Planofix (Alpha NAA) at a very low dose (0.25 ml per liter) to improve fruit set.

			Install Yellow Sticky Traps (10/acre). Spray Imidacloprid (0.3 ml/L) or Neem Oil.
Mango	Fruit development and maturity stage	Fruit fly	Provide light irrigation at 10–12-day intervals if soil moisture is low. Apply Sulphate of Potash (00:00:50) @ 5g/liter of water to improve fruit quality and weight. Apply a foliar spray of Potassium Nitrate (KNO₃) at 1–2% to improve fruit size, sweetness and heat tolerance Install Methyl Eugenol pheromone traps (10 traps/hectare).
Arecanut		Inflorescence Die-back	Apply dry leaves, coconut husk or organic mulch in basins to conserve moisture. Stem Wrapping: Wrap the exposed south-western side of the palm stems with dry areca leaf sheaths, opaque polythene, or trunk with lime (calcium carbonate) . This prevents the stem from heating up, cracking, and eventually breaking. Spray Hexaconazole (1 ml/liter) or 1 % Bordeaux mixture.

Livestock, Poultry, and Sericulture Advisory	
Sector	Weather-Based Advisory
Livestock	<ol style="list-style-type: none"> 1. Provide dry and clean shelter; avoid animals standing in wet areas. 2. Provide clean and cool drinking water 3–4 times/day. Ensure sufficient water availability at all times. 3. Monitor for tick and mite infestations; use approved acaricides if needed. 4. Provide balanced feed and mineral supplements. 5. Provide shade, proper ventilation, and sprinkle water on shed roof if needed during peak afternoon heat. 6. Continue vaccination and deworming schedules. Observe for signs of heat stress (panting, reduced feed intake).
Sericulture	<p>Chawki (early stage): Maintain temperature 26–28°C and RH 75–85%. Sprinkle water on floor to maintain humidity if required.</p> <p>Late age larvae: Maintain temperature 24–26°C and RH 65–70%. Ensure good ventilation to prevent disease incidence.</p> <p>Provide fresh, tender leaves in required quantity. Avoid wilted leaves.</p> <p>Disinfect rearing house and appliances before batch rearing. Remove bed refuse regularly.</p>
Poultry	<ol style="list-style-type: none"> 1. Ensure proper airflow in poultry sheds. Use fans if necessary. 2. Provide cool water with electrolytes during hot hours. 3. Provide electrolytes + vitamins in water for immunity. 4. Keep litter dry to prevent ammonia accumulation. 5. Maintain biosecurity measures and regular health monitoring..

Block level weather forecast (From 06-05-2026 to 10-05-2026)**CHANNAPATTANA BLOCK**

Parameter	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
Rainfall (mm)	4	1	5	1	2
Max. temp (°C)	34	35	35	35	35
Min.Temp (°C)	24	24	25	25	25
Sky condition (Octas)	6	4	5	4	5
Relative humidity (%) 0830 hours	75	76	75	77	77
Relative humidity (%) 1730 hours	36	35	30	31	32
Wind Speed (kmph)	5	10	4	4	3
Wind Direction	288	264	225	158	140

KANAKAPURA BLOCK

Parameter	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
Rainfall (mm)	2	2	6	1	2
Max. temp (°C)	34	34	35	34	34
Min.Temp (°C)	24	24	25	25	25
Sky condition (Octas)	6	5	6	5	6
Relative humidity (%) 0830 hours	77	77	81	82	80
Relative humidity (%) 1730 hours	38	38	33	36	36
Wind Speed (kmph)	4	7	5	4	7
Wind Direction	349	264	162	138	119

MAGADI BLOCK

Parameter	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
Rainfall (mm)	3	1	5	1	4
Max. temp (°C)	33	34	35	35	35
Min.Temp (°C)	23	24	25	25	25
Sky condition (Octas)	6	4	6	4	6
Relative humidity (%) 0830 hours	76	74	72	72	74
Relative humidity (%) 1730 hours	35	35	29	30	30
Wind Speed (kmph)	4	11	7	5	9
Wind Direction	253	262	214	168	137

RAMANAGARA BLOCK

Parameter	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
Rainfall (mm)	4	1	5	1	2
Max. temp (°C)	34	35	35	35	35
Min.Temp (°C)	24	24	25	25	25
Sky condition (Octas)	6	5	5	4	5
Relative humidity (%) 0830 hours	75	75	75	76	77
Relative humidity (%) 1730 hours	36	35	30	31	32

Wind Speed (kmph)	4	10	5	5	4
Wind Direction	292	264	209	157	127

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: mausam.imd.gov.in


For any information farmers can contact **Dr. M. N. Thimmegowda**, Professor & Head/
Mr. L. Nagesha, Technical officer over phone No. **9741109702/ 9008454142**

AMFU of IMD,
AICRP-AM, Bengaluru

वास्तविक वर्षा तथा विस्तारित अवधि पूर्वानुमान
Realized Rainfall and Extended Range Forecast
(वर्षा और तापमान)
(Rainfall and Temperature)

Realized Rainfall

(16th to 29th April 2026)




- Normal or above normal rainfall occurred in both the weeks over Sub Himalayan West Bengal & Sikkim, Assam & Meghalaya and North Interior Karnataka.
- Normal or above normal rainfall occurred in either of the two weeks over Haryana-Chandigarh-Delhi, Bihar, Nagaland-Manipur-Mizoram-Tripura (NMMT), East Madhya Pradesh, Madhya Maharashtra, Marathwada, Konkan & Goa and Coastal Karnataka.
- Below Normal rainfall / No rain occurred in both the weeks over rest of the States & UTs.


Extended Range Forecast System

Rainfall forecast maps for the next 2 weeks (IC- 29th April, 2026) (01st to 14th, May 2026)

INDIA METEOROLOGICAL DEPARTMENT Rainfall Forecast Map - Normal RF Based Categories




INDIA METEOROLOGICAL DEPARTMENT Rainfall Forecast Map - Normal RF Based Categories



- **Week 1 (01.05.2026 to 07.05.2026):** Rainfall is likely to be above normal over North East India, Bihar, West Bengal & Sikkim, East Uttar Pradesh, Kerala and South Interior Karnataka. Rainfall activity is likely over Jammu & Kashmir, Uttarakhand & Himachal Pradesh, over many parts of East India and South India and at isolated places over Madhya Pradesh, Chhattisgarh and East Rajasthan
- **Week 2 (08.05.2026 to 14.05.2026):** Rainfall is likely to be above normal over Kerala, South Interior Karnataka, Coastal Karnataka, Konkan & Goa, Saurashtra-Kutch, Uttarakhand, Bihar and Gangetic West Bengal, over many parts of North East India and some parts of West

Rajasthan. Rainfall activity is likely over many parts of East India, South India, Jammu & Kashmir, Himachal Pradesh, Uttarakhand and Punjab.

**Maximum and Minimum temperature anomaly ($^{\circ}$ C) forecast
for the next 2 weeks (IC – 29th April, 2026)
(01st to 14th, May 2026)**




Maximum Temperature (Tmax)

- **Week 1 (01.05.2026 to 07.05.2026):** Maximum Temperature is likely to be above normal over West Rajasthan and some parts of Saurashtra-Kutch and Odisha. It is likely to be below normal over rest of the country.
- **Week 2 (08.05.2026 to 14.05.2026):** Maximum temperature is likely to be above normal over many parts of North West India, Gujarat State, West Madhya Pradesh, Vidarbha, Konkan & Goa, Chhattisgarh and Odisha.

MME forecast Tmin anomaly (Deg C)

(Week1: 01May-07May)

(Week2: 08May-14May)



Minimum Temperature (Tmin)

- **Week 1 (01.05.2026 to 07.05.2026):** Minimum Temperature is likely to be above normal over Rajasthan, Punjab, Haryana, Gujarat State, Konkan & Goa and many parts of Uttar Pradesh and South India.
- **Week 2 (08.05.2026 to 14.05.2026):** Minimum temperature is likely to be above normal over many parts of South India, North West India, Central India and West India.